110 alo<u>y</u>, 4(9109)

MANADIR LE ANNE RIO DERNA E

PSYCHOLOGY

G STANDEY HALD.

vand Psychiermany, April n of Professor r and Danne-

gne. Address iatry, Univer-

in connection and Psycho-of June, 1909. methods, and d courses by professionally sh to prepare work, a survey cordingly, the

d demonstras and experi-

of the subject be considered.

brain. Child of the feeble-training. Infeebleminded ensic Psychiatics (Articula-and cripples. linics and Scine will appear Privatdozent ankfurt, a. M.,

d:

. M.

SANFORDS AND E. B. TUTCHENER.
LUmparity Cornell University.

WITH THE COTORRECTION OF A

WITH THE CO-ORDER CON-ORDER CONTROL OF CONTR

OVERTHER STREET, WE WEEK !

THE INFLUENCE OF THE DURATION AND OF THE RATE OF ARM MOVEMENTS UPON THE JUDGMENT OF THEIR LENGTH

By PROF. J. H. LEUBA, assisted by MISS ETHEL CHAMBERLAIN, Fellow in Psychology, Bryn Mawr College

Description of the Experiments. During the past year I undertook, with the help of the apparatus described in the preceding pages under the name of "An Apparatus for the Study the duration and of the velocity factors in the estimated the duration and of the velocity factors in the estimation of the length of arm movements made by the forearm. Two of these parts are still unfinished. They deal, one with the source of the constant error arising when a length, objectively limited, is reproduced by a free, i. e., unlimited, movement; the other with the effect of voluntary changes in the velocity with which the reproducing movement is made.

The part here described consists of four series: 1. a Normal Series (N. Ser.)¹; 2. a Weight Resistance Series (W. R. Ser.)¹; 3. a Muscle Resistance Series (M. R. Ser.)¹; 4. a Control Series, or an Increasing Weight Resistance Series (I. W. R.

The method used throughout was the modification of the Method of Average Error called the Method of Constant Stim-The observer's task in this method is to compare a standard movement with a number of fixed comparison-lengths so chosen that the smallest seems distinctly shorter and the largest clearly larger than the standard (St.)1, and to express his judgment by one of the words, "smaller", "equal" "greater". I have used only three comparison lengths (C.)1. The results show this small number to have been sufficient for our purpose.

In each of the four series, the St. Movement (Mt. in the singular and Mts. in the plural) was a length of 8° gone through in the upward direction and beginning each time at a different point between 0° and 5°. The zero point is the one indicated by the pointer when the arm is held horizontally. The reason for this latitude in the starting point of the St., and for a similar latitude in the starting point of the C. was

the wish to make it impossible.

In the N. and in 25° and 30°, i. e., a of the St. in all Ser.

In the N. Ser. the a resistance of 11/2 weight attached to a rim of one of the wh weight was put on o

In the M. R. Ser. semi-circle as to end from the extreme pos was provided in this subject so that his po remain constant. A pressure of the forea duced. It is to be no from the beginning to in the W. R. Ser. conclusion.

Two subjects perfor graduate student in who was the only subj the experiment and th (H.), a senior studen experience in the com R. and the M. R. Ser.

The observers worked kept them closed withc before each new Ser. as session. These session duration, broken by set

Introspection was ask series.

The directions given concentrate their attent express by the words, " relation of the C. to the locity" were not mentio trospective account was move at the speed they l

Results. I. I give fin the N. Ser. of L. (Table fiftieths of a second, the when the C. were respect judgment passed by the

¹These abbreviations will be used throughout the article.

DY OF KINÆS-TION.

ollege, Pa.

he apparatus as set

ver the lever (L.).
distance from the
t is opposite the end
on the lever, but is
holder, and at the
hand of the subject
instead of making
tied to the lever
When the point of
oper arm hangs vers horizontal.


paratus are the fol-

rotation of a shaft ibed by the moving ion of the arm, i. e.,

metallic blades at ades are connected with an ordinary rected a spring (sp) us an electric circuit passes under the

be made to move at cond, several wheels cted with a separate between the blades movements of any × 300), can be re-

cis so that they proneously. ve downward movements made by each one of the recording magnets corresponds to an arm movement of 2°. But by combining the lines traced



the wish to make judgments by means of the sense of position impossible.

In the N. and in the W. R. Ser., the C. Mts. began between 25° and 30°, i. e., approximately midway between the position of the St. in all Ser. and that of the C. in the M. R. Ser.

In the N. Ser. the Mt. was unimpeded; in the W. R. Ser. a resistance of 1½ kilogr. was provided in the form of a weight attached to a wire running in a groove made in the rim of one of the wheels, the whole being so disposed that the weight was put on only after the St. and before the C. Mt.

In the M. R. Ser., the C. lengths were so placed on the semi-circle as to end at different points within 3° of a point 1° from the extreme position the arm was able to reach. A rest was provided in this series for the back and the head of the subject so that his position with regard to the apparatus would remain constant. A very considerable resistance, due to the pressure of the forearm against the biceps, was thus introduced. It is to be noted that this resistance was not constant from the beginning to the end of the C. Mt., as it was the case in the W. R. Ser., but that it increased steadily until its conclusion.

Two subjects performed the four Ser., one of them (R.), a graduate student in psychology, the other (L.), the writer who was the only subject informed regarding the purpose of the experiment and the detail of procedure. A third subject (H.), a senior student, who had already had a good deal of experience in the comparison of arm Mts., served for the W. R. and the M. R. Ser.

The observers worked with their eyes bandaged, except L. who kept them closed without that help. Practice series were given before each new Ser. and a few tests at the beginning of every session. These sessions were limited to about half an hour's duration, broken by several interruptions for rest.

Introspection was asked for only towards the end of each

Series.

The directions given to the subjects were simply that they concentrate their attention upon the length of the Mts. and express by the words, "shorter", "equal", and "longer", the relation of the C. to the St. The words "duration" and "velocity" were not mentioned to them, at least not before the introspective account was asked for. They were thus left to move at the speed they liked best.

Results. I. I give first, as a sample, a complete record of the N. Ser. of L. (Table I). The figures indicate, in terms of fiftieths of a second, the duration of the St. and of the C. Mts. when the C. were respectively 7°, 8°.5, and 10° and also the judgment passed by the subject in each case. Observe the

TABLE I
Normal Series

00			

Subject L Equality judgm	1
Equality indea	103
- Tanasay Judgm	ent m

C. = 7°						C = 8°.5				C. = 10°.					1			
			==	= 	+				=		+				=		+	
	St.	c.	St.	c.	St.	c.	St.	c.	St.	c.	St.	c.	St.	c.	St.	c.	St.	0
	67 84 63 63 60 83 60 75 83 79 74 67 90 71 88 87 74	85728 828 828 838 849 849 854 854 854 854 854 854 854 854	83	88 72 68 42 48 81			73 74 78 63 79 73 79 69	80 58 62 68 56 58 77 83	75 81 77 94 86 75 75 95 85 76 47 55 63 60 106	76 77 93 69 82 65 75 69 64 55 52	822 800 67	87 91 88			83 71 111 86 58	101 97 88 88 66	79 73 88 86 75 73 73 70 70 70 70 71 79 70 70	* FINDERFERENCE IN REPORT
Sums	1601	1221					588	542	1231	1210							1664	195
Aver.	76	61					73 · 5	68	77	76							76	Si

considerable variable error in the duration both of the St. and of the C. In column 7°—, for instance, the St. times vary from 58 to 92. The differences between the compared St. and C. are also considerable. They range in the same column from +19 to -40. Similar variations were found in series of experiments in which the observer was comparing not the lengths of Mts., but the *duration* of pressures upon his forefinger. I shall use this similarity in a subsequent article to reinforce the thesis that in the estimation of the length of Mts. here dealt with, it is the duration and the velocity, not the length, which are directly perceived. Nevertheless, the average duration of the St., leaving out the columns containing

too few test column 8° uniformity tests taker

The dur series fron He moved

The mc ing the diand felt eidentical tits duratic 68). If vof 10° (felonger tin ing in this is practic 76+0.5

siderably columns (variable ! These:

These: to be rein thinks, to two move add that mind. I movemen his attent joint.

II. In the sum columns the num difference renthesis of these number (marked others. in which the semi however

If the appeared

too few tests, is strikingly uniform: 76 for column 7°—; 77 for column 8°.5—, and 76 for column 10°+. We look upon this uniformity as an indication of the sufficiency of the number of tests taken.

The duration of the individual Mts. varies for L. in this series from a little below one second to a little over two seconds. He moved more rapidly in the M. R. and the I. W. R. Ser.

The most interesting figures of the table are those expressing the duration of the C. When the C. was $8^{\circ}.5$ in length and felt equal to the St., St. and C. were made practically in identical times (77 to 76); when the same C. was felt shorter, its duration was materially less than that of the St. (73.5 to 68). If we pass to the comparison of the St. with a C. length of 10° (felt markedly longer), we find that it took a much longer time to make the C (76 to 89). The proportion existing in this case between the duration of the compared movements is practically equal to that existing between their lengths: $10^{\circ} + 10^{\circ} = 10^{\circ}$ When the C. Mt. was 10° it was made in considerable that there there the St. (76 and 61). The other

siderably less time than the St. (76 and 61). The other columns contain too few tests to yield an average free from the variable time error to which I have drawn attention.

These figures lead, it seems, to the conclusion—a conclusion to be reinforced by our other results—that in comparing, as he thinks, two lengths, L. compares in reality, the duration of two movements made at approximately the same rate. I may add that during the experiment he did not have duration in mind. He strove to reproduce in the C. the sensations of movement experienced in the St. The sensations that drew his attention seemed to be localized in and around the elbow joint.

II. In order to get the duration figures given in Table II, the sums (not the averages) of the —, the —, and the + columns in each series were added, and the total divided by the number of tests included in each column. The differences between the St. and the C. are indicated in parenthesis. The W. R. Ser. of H. includes but 50 tests; of these only ten fall in the + column, hardly a sufficient number for a reliable average. The figure in the equality C. (marked with an asterisk) is not strictly comparable with the others. It is derived from a series not reported in this paper in which the C. Mts. were made considerably higher up on the semi-circle than in the N. Ser. The figure may serve, however, as an approximation.

If the judgments recorded in this table are based, as it appeared probable from the series already examined, upon time

appeared pro

JOURNAL---6

= 10°.

idgment 8°.4

664 1967

76 89

and vary and umn es of

the forele to Mts.

ver-

TABLE II

Average duration of all the tests arranged by series and classified according as the C. was judged larger, equal, or smaller than the St.

Subject	Series		_			=			+		lity
Sul	Se	St.	Diff.	С.	St.	Diff.	C.	St.	Diff.	C.	Equality C.
	N	75	(—14)	61	79	(76		(+14)	90	8°.4
Ļ	WR	76	(—ro)	66		(+ 1)	79	80	(+11)	91	80.3
	MR		nough cases.	(6)	48	(- 3)	45	54	(+ 2)	56	7°.3
	IWR	50	(- 1)	49	51	(o)	51	50	(+ 9)	59	7°.9
	N	39	(+ 2)	41	37	(+ 5)	42	37	(+ 9)	46	8.7
R	WR	41	(+ 8)	49	40	(+11)	51	41	(+19)	60	8.5
1	MR	32	(+ 6)	38	31	(+ 8)	39	31	(+14)	45	7.5
	IWR	43	(+ 5)	48	40	(+ 7)	47	40	(+13)	53	7.6
	N										10*
H	WR	30	(+ 6)	36	29	(+11)	40	31 (+	- IO) IO cases.	40	9.8
	MR	29	(+11)	40	29	(+15)	44		(+16)	47	7.2

comparisons, we should expect the duration of the compared Mts., in every case in which they are judged of equal length, to bear to each other a fixed relation, whatever be the length of the C. That fixed relation would be one of equality if the subject moved through the C. with the same velocity as through the St. But if he should move systematically slower through one of them, the constant relation could not be one of equality. I assume, of course, in saying this, that the subject is influenced, not only by the duration but also by the speed of his Mts., that, in fact, a quasi-automatic compensation takes place between duration and speed when he is judging of spatial lengths.

The data before us bear out these suppositions. Subject L., moving through the St. and the C. in the N. and in the W. R. Ser. at an approximately equal velocity, finds the C. equal to the St. when both take the same time, shorter than the St. when the C. takes less time, and longer than the St. when it is the reverse. In the M. R. and the I. W. R. Ser., done several days after the others, he again maintains practically the same rate of speed through the St. and the C. Mts., although the absolute speed is considerably increased when compared with that of the first two Ser. Subject R. differs from L. in that she takes the C. more deliberately than the St., and, allowing for the lesser speed of the C. Mts. calls, in the first series, the C. equal to the St. when the former takes her 5 units longer. When it takes her only 2 units longer, she calls it shorter, and

when it lasts 9
Ser. she procee
preceding Ser.,
not clearly con
Her Equality (
the C. takes he
when it lasts I
Her last two S
her first two S
ceed from a si
the two other s

We find, the to the task of were guided in the Mts., alth awareness of changes of ab St. and C. a prise. And the subjects did now as, I confess at some future length of two deavor to move

Explanation the average 1 St. of 8°. tical column to the influe every subject i. e., the equ differ materia looked that it not alter the in the M. R. tial overestir tion of the (of Table II a shorter tim the lengths If no times. tion of the ra R. Ser. a gr arm moves t the increasing pressing age eccord-

8°·4 8°·3 7°·3 7°·9 8·5 7·6 10* 9.8

7.2

red th, a of ub-

igh ıgh ity.

fluhis kes tial

St. t is ral me :he ith

W. ual

ıat ng :he er. nd. when it lasts 9 units more, she calls it longer. In the W. R. Ser. she proceeds through the C. still more slowly than in the preceding Ser., and the figures show that, even though she is not clearly conscious of it, this decrease is taken into account. Her Equality C. takes her 11 units longer than the St. When the C. takes her only 8 units longer, she calls it shorter, and when it lasts 19 units more than the St., she calls it longer. Her last two Ser. are, in this respect, in full agreement with her first two Ser. The averages from H., as far as they proceed from a sufficiently large number of tests, match those of the two other subjects.

We find, thus, conclusive evidence that these subjects, set to the task of comparing the length of two arc movements, were guided in their judgments by the duration and the rate of the Mts., although their introspection does not indicate any awareness of the fact. The degree of precision with which changes of absolute speed and of relative speed between the St. and C. are taken into account, came to me as a surprise. And the further fact that, making use of duration, the subjects did not try to keep the velocity equal (L. excepted) was, I confess, somewhat disconcerting. I intend to find out at some future time whether persons required to compare the length of two Mts. by means of their duration would not endeavor to move with an equal speed through both.

Explanation. Let us, now, compare in the different series the average lengths of the C. when it was felt equal to the St. of 8°. These lengths are written down in the last vertical column under the head "Equality C." With regard to the influence of the weight resistance, they confirm, in every subject, the results obtained by other experimenters, i.e., the equality C. in the N. and in the W. R. Ser. do not differ materially one from the other. It should not be overlooked that it is the addition of a constant resistance which does not alter the length of the Mt. The increasing resistance used in the M. R. and in the I. W. R. Ser. produced a substan-But why this overestimatial overestimation of the C. Mts. tion of the C. in the M. R. Ser.? We know from the figures of Table II that it has not its origin in an error of duration: a shorter time is not mistakenly judged equal to a longer one; the lengths 7°.3 and 8° felt equal by L. are made in equal times. If not in the duration, the error must be in the estimation of the rate of Mt. There is as a matter of fact in the M. R. Ser. a gradual reduction in the velocity with which the arm moves through the C. This falling off is correlated with the increasing resistance offered to the Mt. by the forearm pressing against the upper arm. We shall submit figures on this point in a future paper. But the falling off of the rate is not in itself sufficient to cause an error since, as we have seen, into account. It must be that in this Ser. the conditions do not allow of the normal compensatory effect of the speed upon

At this point we must inquire into the sensory basis for the apprehension of the rate of Mt. The skin sensations,—air pressure and temperature—at best secondary criteria, had been eliminated in the cases of R. and H., by means of a loose fitting glove, reaching almost to the elbow, and overlapped by the sleeve. These sensations being out of the question, there remained those arising from the muscles, from the tendons, and from the joint. When the arm is moved through a fixed number of degrees, the effort made is greater when the velocity is high than when it is low, because a part of the innervation finds its way into the antagonistic muscles, and an increased innervation of the muscles that do the work is required. But I do not see any reason for assuming that, as far as tendinous and muscle sensations are concerned, the untrammelled arm moving upward at a considerable speed yields a sensation-experience different from that produced by a slower movement of the same duration made against a properly chosen resistance. The muscle activity would, it seems, be the same in both cases. The muscle and tendinous sensations cannot, therefore, provide an unequivocal basis for the rate of Mt. One is, thus, driven to the hypothesis that unequivocal speed information comes from the joint surfaces.

What is it, then, that takes place in the joint surfaces when speed changes? Two things: (1), an increase in the pressure of the joint surfaces against each other, corresponding to the greater innervation of the muscles. This pressure-increase produces, presumably, a rise in the intensity of the joint sensations. But an intensity series correlated with an increase in muscular tension cannot, as we have seen, serve as a basis for an apprehension of speed. (2.) Speed-increase means also an increment in the rapidity with which the joint surfaces pass over each other, i. e., an increase, shall I say, in the quantity of the stimulus applied to each one of the joint senseorgans in a unit of time? It is credible enough that a qualitative or a quantitative sensation-difference corresponds to this quantitative increase of the stimulus. We may think of it as being the same sort of difference as the one experienced when the velocity of a wheel, the edge of which rubs against the skin, is altered. Whatever be the psychical effect produced, in the case of the joint organs, by changes in the rate with which the stimuli succeed each other, that seems clearly

the only possible equivocally to the I shall therefore hand, that our comparison of during upon the rap stimulated. This the joint sensation year the existence

effective presence We are now pr the rate of Mt. in the length errors that as the C. r somewhat irregula not for the physic the effort made at creased resistance of its purpose, sin resistance has aga not of itself caus plicating circumst rate impossible th is that the rate-va gradual and some joint sensations. muscular tension Furthermore, and ments of the spec tion away from th The error is an c

speed is greater the In the W. R. S tioned did not ex muscle tension in sponding increme But this increase in ning to the end c divert the attention error took place.

If the duration of Ser., it is, I think as a whole, or of t mated by means of unaffected by the ent matter whether

ig off of the rate is ce, as we have seen, itions, exactly taken in the conditions do ct of the speed upon

sensory basis for the kin sensations,—air idary criteria, had H., by means of a ie elbow, and overbeing out of the m the muscles, from the arm is moved fort made is greater low, because a part ntagonistic muscles, uscles that do the reason for assuming tions are concerned, a considerable speed n that produced by nade against a propity would, it seems, nd tendinous sensavocal basis for the sypothesis that unejoint surfaces.

joint surfaces when case in the pressure corresponding to the is pressure-increase ity of the joint senwith an increase in serve as a basis for crease means also an joint surfaces pass [say, in the quanof the joint sensemough that a qualince corresponds to

We may think of he one experienced which rubs against sychical effect prochanges in the rate that seems clearly the only possible cause of a sensory change corresponding unequivocally to the rate of the Mt.

I shall therefore hold, until further information comes to hand, that our comparative judgments of the length of arm Mts., when the "sense of position" is excluded, is really a comparison of duration and of a peculiar sensory value depending upon the rapidity with which successive joint organs are stimulated. This sensory value may be called the *rate-value* of the joint sensation. That introspection does not clearly reveal the existence of this rate-value is no argument against its

We are now prepared to account for the over-estimation of effective presence. the rate of Mt. in the M. R. Ser, an over-estimation leading to the length errors recorded in Table II. I have said already that as the C. movement proceeds the resistance increases somewhat irregularly and the velocity falls off. It decreases, not for the physiological reason offered by Loeb, but because the effort made at any particular moment to overcome the increased resistance, so as to keep the speed constant, falls short of its purpose, since by the time the effort becomes effective the resistance has again increased. The decrease in speed would not of itself cause the observed error. It is because of complicating circumstances making the correct estimation of the rate impossible that the Mt. is made shorter. What happens is that the rate-value of the joint sensation is obscured by a gradual and somewhat irregular increase in the intensity of the joint sensations,—an increase arising from the increment in muscular tension made necessary by the growing resistance. Furthermore, and chiefly, the necessity of repeated readjustments of the speed to the resistance tends to draw the attention away from the rate-value.

The error is an overestimation of the rate because the intended

speed is greater than the realized one.

In the W. R. Ser. the disturbing circumstances just mentioned did not exist. There was, indeed, an increase of the muscle tension in the C. Mts., and, presumably, a slight corresponding increment in the intensity of the joint sensations. Sponding increase was, in this series, constant from the begin-but this increase was, in this series, constant from the beginning to the end of the C. movement. It did not, therefore, ning to the attention from the rate-value and thus no velocity divert the attention from the rate-value and thus

error took place.

If the duration of the Mt. remains undisturbed in the M. R.

Ser., it is, I think, because the duration of the Mt. experience
sas a whole, or of the joint sensations in their entirety, is estias a whole, or of the duration of other sensations themselves
mated by means of the duration of other sensations themselves
unaffected by the arm movement. It is, therefore, an indifferunaffected by the arm movement any particular qualitative or inten-

sive alteration takes place in any one of the sensory components of the whole Mt. experience. The important points for a correct duration-estimate are that the Mt. should be present in some way in consciousness and that the sensations used to measure its duration—and we know that they are not necessarily the same for every one—remain undisturbed by extraneous factors as long as the Mt. lasts.

In order to verify the preceding explanation of the space error present in the M. R. Ser., a control Ser. was devised, In it the C. Mts. took place over the same portion of the semicircle as in the W. R. and the N. series. A resistance was introduced in such a way that, instead of being constant as in the W. R. Ser., it increased as the C. Mt. proceeded. Thus the pressure of the forearm against the upper arm, and the high degree of contraction of the biceps were eliminated, but the increasing resistance remained. Under these circumstances one would expect, if our explanation is valid, a rate error just as in the M. R. Ser. The resistance provided was a long weight of 2 Kgr. dipping in mercury. During the C. Mt. the weight was gradually lifted out of the mercury. As, according to our practice, the starting point of the C. as well as of the St. varied by a few degrees in this series also, the initial resistance in the C. Mt. varied from o to about 300 grams, The amount of resistance at the end of the Mt. was not at all as great as in the M. R. Ser.; nevertheless, the results show, in the case of R., an overestimation of the equality C. practically equal to the one present in the M. R. Ser. (7°.6 against 7°.5 in the M. R. Ser.). The error is not so great in the case of L. (7°.9 against 7°.3 in the M. R. Ser.), yet an unmistakable error in the same direction as in the M. R. Ser. is apparent in his results. It is in fact the size of R.'s error and not that of L. which is surprising in view of the considerably smaller resistance applied in this Control Ser. Lack of time prevented H. from doing this series.

Our account of the over-estimation of the equality C. is thus verified, as far as it was in the power of the Control Series to do. Historical and Critical. In 1890 J. Loeb published in Vol. 46 of Pflüger's Archiv, pp. 1-46, under the title, "Untersuchungen über die Orientirung im Fühlraum der Hand und in Blickraum", certain experiments with arm movements from which he concluded that the more contracted are the active muscles at the beginning of a movement, the greater is the

muscles at the beginning of a movement, the greater is the overestimation (p. 41). In order to account for this fact he makes use of two hypotheses, the first of which is superfluous, and the second (dependent upon the innervation theory) is now quite discredited. They are, (1.) the excitability of a muscle decreases as its state of contraction increases; (2.) the

length of a Mt. muscles perform Mts., the second and is therefore of contraction, of the second amount by who contracted muones, an equal in the second to

F. Kramer den Lage- u Psy., Vol. XX ments and, go movements in maining equa of the other. gested, witho test, that the come of a nat difficult Mt. statement we resistance, a tended mean.

The assum Loeb's illusic ''Ueber die empfindunge He used for constructed paper it rec through a ri a record is r In one of th with the ha a certain dis same directi first. In th the observe His dur

may take is the times of the contact Loeb's e

the sensory components portant points for a cor-It. should be present in the sensations used to hat they are not necesindisturbed by extrane-

cplanation of the space ntrol Ser. was devised. ame portion of the semi-A resistance was :ies. of being constant as in Mt. proceeded. Thus the upper arm, and the ps were eliminated, but ider these circumstances s valid, a rate error just e provided was a long During the C. Mt. the mercury. As, accordof the C. as well as of s series also, the initial o to about 300 grams. the Mt. was not at all ieless, the results show, f the equality C. practi-1. R. Ser. (7°.6 against not so great in the case Ser.), yet an unmistakthe M. R. Ser. is appare of R.'s error and not ew of the considerably trol Ser. Lack of time

of the equality C. is thus the Control Series to do. Loeb published in Vol. er the title, "Untersuchraum der Hand und in i arm movements from ntracted are the active nent, the greater is the account for this fact he of which is superfluous, innervation theory) is (a) the excitability of a ction increases; (2.) the

length of a Mt. is judged by the amount of energy sent to the muscles performing it. If, then, we try to make two equal Mts., the second of which follows in the direction of the first and is therefore performed with the muscles in a greater state of contraction, we shall send to the muscles for the execution of the second movement an amount of energy equal to the amount by which the first was produced. But as the more contracted muscles are less excitable than the less contracted ones, an equal amount of energy will produce a shorter length

in the second than in the first Mt.

F. Kramer and G. Moskiewicz in "Beiträge zur Lehre von den Lage- und Bewegungsempfindungen' in the Zeits. f. Psy., Vol. XXV (1901), pp. 101-125, repeated Loeb's experiments and, generalizing his conclusions, stated that of two movements intended to be of equal length, everything else remaining equal, the more uncomfortable (unbequem) falls short of the other. They rejected Loeb's explanation, and suggested, without putting their supposition to an experimental test, that the ovestimation of the "unbequem" Mts. is the outcome of a natural tendency to move more slowly through a difficult Mt. than through an easy one (pp. 121-123). statement we know to be inexact, unless the effect of an added resistance, as in our W. R. Ser. does not fall within the intended meaning of the word "unbequem".

The assumption of Kramer and Moskiewicz with regard to Loeb's illusion was recently put to a test by Erich Jaensch in "Ueber die Beziehungen von Zeitschätzung und Bewegungsempfindungen", Zeits. f. Psy., Vol. 41 (1906), pp. 257-279. He used for measuring the duration of the Mts. a pencil so constructed that when its point is pressed down upon the paper it recedes into the handle. Air is thus pressed back through a rubber tube into a Marey's tambour. In this way a record is made of the beginning and of the end of the Mts. In one of these two series of experiments the observer, starting with the hand near the chest, moved it away from the body for a certain distance, stopped an instant, and then proceeded in the same direction until he had made a line seemingly equal to the first. In the other set, the direction of the Mts. was reversed, the observer starting away from the body and coming toward it.

His duration records led him to this conclusion: "One may take it as proven that the lengths appear equal because the times used in making them are equal" (pp. 269). He discusses the causes of the lower velocity of the uncomfortable Mts. and comes to the opinion, different from ours, that the chief one is physiological, namely, the decreasing excitability of the contracting muscle.

Loeb's explanation of the decrease in length of the Mt. is

thus taken up by Jaensch as the cause of the decrease in its rapidity. This decreasing excitability hypothesis is shown by our Increasing Weight R. Ser. to be superfluous. Moreover, if the relation it supposes between the degree of contraction and excitability really existed, we would in all likelihood have learned to increase automatically the innervation as the length of the muscle decreases and thus have overcome the tendency to error. Taensch does not seem to have clearly realized, any more than his predecessors, what our experiments prove, namely, that it is an increasing resistance and not an added, constant resistance which causes the velocity error in Loeb's experiments. Neither did he grapple with the real problem, which was to make clear why, while we are usually able to take into account, for the sake of the length of the Mt., minute changes of velocity, under special conditions, a constant error in the estimation of speed takes place. To this problem an answer has been given in the preceding pages.

Introspection revealed that our three subjects made no use of visual imagery. In the case of L, there was at times a vague image of the space covered, but it followed the production of the Mt. and never had the clearness necessary in order to serve as a guide. One of the two subjects we used in other experimental series, and only one, had clear visual images of the length of the Mt. She saw the St. and tried to reproduce an equal visual length. Of all our subjects she was the least accurate. I hope to have the opportunity of finding out whether she also made use of duration and rate in judging of the length of Mts.

In the light of the preceding facts and discussions the complete inadequacy of Külpe's statement is evident: "our judgment of the extent of arm movement is not based (in the author's observation) upon the temporal relations of the movement, but upon the reproduced visual image of the space passed through, and more especially of the extreme positions

of the moved arm." (Outlines, p. 348.)

Before closing I must refer to the constant positive error in the Equality C. of the N. and of the W. R. Ser. in the case of the three subjects. (See the last column of Table II.) I have so far neglected it. My reason for doing so is that I have at present nothing decisive to say about it, I do not know how to account for the overestimation of the velocity, that, in my view, it implies. There are several possibilities. I shall probably return to the point in another paper.

Summary of Conclusions. When the sense of position is ex-

cluded, the comp through the comp sensations arising tions) and of a here the rate val

A quasi-auton

A gradual in whether caused against the uppe Mt. This decrestimated, and movement takes

Local signatu joint sensations, of a whole range whole, not the local sign.

The comparis present in our e of local signs i duration and of

decrease in its sis is shown by Moreover, if traction and exod have learned e length of the ndency to error. any more than e, namely, that ed, constant re-)'s experiments. 1, which was to o take into acinute changes of error in the estian answer has

ects made no use was at times a wed the producecessary in order we used in other r visual images d tried to reproects she was the y of finding out te in judging of

ussions the coment: ''our judgot based (in the elations of the sage of the space xtreme positions

positive error in er. in the case of of Table II.) I so is that I have do not know how city, that, in my ies. I shall pro-

of position is ex-

cluded, the comparison of the length of arc movements is made through the comparison of the duration of one or several of the sensations arising from the Mts. (preferably the joint sensations) and of a particular value of the joint sensation, called here the rate value.

A quasi-automatic compensatory relation exists between the

duration and the rate value.

A gradual increase in the resistance offered to a Mt., whether caused by a weight or by the pressure of the forearm against the upper arm, produces a decrease in the rapidity of the Mt. This decrease, for reasons mentioned above, is underestimated, and thus an overestimation of the length of the movement takes place.

Local signatures cannot be connected with the rate value of joint sensations, for one and the same joint organ is susceptible of a whole range of rate values. It is the joint sensation as a whole, not the rate value, which possesses, or may possess, a

local sign.

The comparison of the length of Mts. under the conditions present in our experiments does not necessitate the existence of local signs in the joint sensations. An apprehension of duration and of rate is sufficient.