



Published in final edited form as:

J Vestib Res. 2018 ; 28(3-4): 283–294. doi:10.3233/VES-180636.

Perception of Threshold-level Whole-body Motion During Mechanical Mastoid Vibration

Rakshatha Kabbaligere^{a,b}, Charles S. Layne^{a,b,c}, Faisal Karmali^{d,e}

^aDepartment of Health and Human Performance, University of Houston, Houston, Texas, United States of America

^bCenter for Neuromotor and Biomechanics Research, University of Houston, Houston, Texas, United States of America

^cCenter for Neuro-Engineering and Cognitive Science, University of Houston, Houston, Texas, United States of America

^dJenks Vestibular Physiology Laboratory, Massachusetts Eye and Ear Infirmary, Boston, Massachusetts, United States of America

^eDepartment of Otolaryngology, Harvard Medical School, Boston, Massachusetts, United States of America

Abstract

BACKGROUND: Vibration applied on the mastoid has been shown to be an excitatory stimulus to the vestibular receptors, but its effect on vestibular perception is unknown.

OBJECTIVE: Determine whether mastoid vibration affects yaw rotation perception using a self-motion perceptual direction-recognition task.

METHODS: We used continuous, bilateral, mechanical mastoid vibration using a stimulus with frequency content between 1 and 500 Hz. Vestibular perception of 10 healthy adults ($M \pm S.D. = 34.3 \pm 12$ years old) was tested with and without vibration. Subjects repeatedly reported the perceived direction of threshold-level yaw rotations administered at 1 Hz by a motorized platform. A cumulative Gaussian distribution function was fit to subjects' responses, which was described by two parameters: bias and threshold. Bias was defined as the mean of the Gaussian distribution, and equal to the motion perceived on average when exposed to null stimuli. Threshold was defined as the standard deviation of the distribution and corresponded to the stimulus the subject could reliably perceive.

RESULTS: The results show that mastoid vibration may reduce bias, although two statistical tests yield different conclusions. There was no evidence that yaw rotation thresholds were affected.

CONCLUSIONS: Bilateral mastoid vibration may reduce left-right asymmetry in motion perception.

1. Introduction

The vestibular system, located in the inner ear, senses three-dimensional angular and linear motion of the head, and provides critical cues for gaze stability, posture and perception. Mechanical vibration applied to the mastoid is known to stimulate the vestibular system. For example, a common clinical “vibration test” for unilateral vestibular hypofunction involves the application of mechanical vibration to the mastoid [42, 46, 92], with the elicitation of nystagmus in the presence of bilateral mastoid vibration (in the range of 60 Hz to 100 Hz) indicating unilateral peripheral dysfunction. Vibration applied on the mastoid has been shown to be an excitatory stimulus to the vestibular receptors [19, 20, 90, 93], using both sinusoidal stimuli (i.e., “pure tones”) ranging from 50 Hz to 4000 Hz and short impulsive stimuli (i.e., “clicks”). It has been proposed that when vibration is applied on the mastoid, the compressional and shear waves generated by the vibration are propagated through the skull towards the inner ear [18, 20]. Upon reaching the inner ear, these bone-conducted vibrational waves displace the receptor hair cells, similar to that produced by natural vestibular stimulation such as angular or linear head motion. In the discussion section we elaborate on the effects of vibration on the two primary vestibular organs, the semicircular canals and the otolith organs.

More broadly, the addition of noise to sensory systems – i.e., a stimulus that does not convey information about the physical quantity of interest – has been shown to influence both sensorimotor and perceptual responses. For example, balance performance changes when electrical noise is applied to the vestibular system [37, 71, 72, 84], the ankle [62] and knee muscles [39], or when mechanical vibration is applied to the bottom of the feet [15, 21, 22, 77] and to the muscle tendons [7]. Similarly, perception of low amplitude visual, auditory and somatosensory signals is modified by the presence of external noise [14, 79, 83, 89, 94]. External noise can either increase or decrease detection or recognition thresholds depending on the level of noise [1, 69]. For example, recent work found that galvanic vestibular stimulation (GVS) with optimal levels of noise lowers perceptual motion thresholds for motion along some axes [33, 34, 52].

Vestibular perception has been studied using a self-motion perceptual direction recognition task in which subject have to report the perceived direction of threshold-level motion stimuli in the dark and their responses are fit using a cumulative Gaussian distribution function that is characterized by two parameters: threshold and bias [10, 11, 17, 49, 87, 97]. Threshold is defined as the standard deviation of the cumulative distribution and refers to the smallest motion that can be reliably perceived. Thresholds quantify the precision of vestibular perception. Bias is defined as the mean of the distribution and refers to the motion perceived on an average when exposed to null stimuli [41, 67]. Vestibular bias may arise from unequal contributions of the left and right vestibular organs or other central imbalances in the sensation and perception of different motion directions [35, 67, 74]. Threshold and bias have been characterized for different planes of motion (e.g., translation along earth-horizontal inter-aural axis, translation along earth-vertical axis, yaw and roll rotation) and with different stimulus frequencies [87]. Studies have demonstrated that proprioceptive, visual or tactile cues from the motion platform are either absent or minimal and that vestibular cues predominate for the perception of motion direction in this kind of task [11, 87]. For example,

patients having bilateral vestibular dysfunction had significantly higher perceptual thresholds when compared to subjects with intact vestibular function [87].

Despite these studies, to our knowledge, the effect of mechanical mastoid vibration on vestibular perception has not been investigated. Hence we tested the effect of vibration on vestibular perception of yaw rotations in a direction-recognition task. For the remainder of this paper, we use “vibration” to refer to continuous, bilateral, mechanical vibration of the mastoids using a random, noisy stimulus with a broad range of frequencies (1–500 Hz). These frequencies were chosen because they correspond to the behavioral bandwidth of vestibulo-ocular responses, which responds up to 50 Hz [44, 78] and even higher [81, 82] and because vibration at these frequencies have been shown to elicit nystagmus and body sway in humans [13, 20, 24, 45, 92]. Recent reports have shown that perceptual vestibular thresholds are correlated with vestibular migraine [57], age [5], peripheral deficits [16, 87], medication use [23] and performance in a clinical balance test [5, 47]. Furthermore, these perceptual thresholds assay neural noise [40], which models have shown impacts dynamic processing of vestibular cues [6, 50, 51, 53, 61]. Thus an improvement in vestibular perception with the use of mechanical vibration may have potential benefits for certain populations.

2. Methods

Ten healthy adults ($M \pm S.D. = 34.3 \pm 12$ years old) participated in the study. The participants were screened using a questionnaire to assess for any history of dizziness, vertigo, back or neck problems, neurological, cardiological or any other general physical problems. The study was approved by the institutional committee for protection of human subjects (CPHS) at the University of Houston and the Institutional review board at Massachusetts Eye and Ear Infirmary (MEEI). All the participants provided informed consent prior to their participation and all study procedures were conducted in accordance with the Helsinki declaration.

All the participants' yaw rotation vestibular perception thresholds and biases were tested under two conditions, with vibration and without vibration. The order of the two conditions was randomized across the participants. Yaw rotation was chosen because it primarily stimulates the semicircular canals, and vibration has been shown to activate semicircular canal neuronal units [93].

2.1 Vibratory stimuli

Vibration was applied bilaterally [13] on the mastoid using two small circular-shaped tactors (C-2 tactor, Engineering Acoustics, Inc.) having a diameter of 1.17 inches and a thickness of 0.3 inches. The tactors were attached to the mastoids using two-sided tape and Velcro straps. The tactors were controlled using custom designed software to generate longitudinal vibrations perpendicular to their surface. These vibrations were band-passed (1–500 Hz, colored noise) and had maximum amplitude of 0.635mm [21, 22]. As no data exists in the literature relating vibratory noise to human vestibular perception, we chose a random, noisy stimulus with a broad range of frequencies to maximize the chance of eliciting a response.

2.2 Motion stimuli

Motion stimuli consisted of single cycles of sinusoidal angular acceleration about the earth-vertical axis at a frequency of 1 Hz. This motion and frequency is known to primarily stimulate the semicircular canals. The stimuli were generated using a motion platform having six degrees of freedom (MOOG CSA Engineering, Mountain View CA, Model 6DOF2000E). The instantaneous angular acceleration of the motion stimulus can be represented by the expression $a(t) = A \sin(2\pi ft)$, where A is the amplitude of acceleration and f is the frequency or the inverse of the period of the stimulus ($T=1/f$). The frequency of the motion stimulus was fixed at 1 second for all trials, which concentrates most energy around 1 Hz [5]. The expressions defining velocity and displacement can be obtained by integrating acceleration and velocity respectively, velocity $v(t) = A/(2\pi f)[1 - \cos(2\pi ft)]$ and displacement $p(t) = A/(2\pi f)[t - 1/(2\pi f) \sin(2\pi ft)]$.

2.3 Experimental procedures

The experimental procedures mimicked those used in earlier studies [11, 38, 49, 80, 87]. All participants were tested in four blocks of 200 motion trials per block. Subjects received mastoid vibration on two of the four blocks. Thus there were a total of 400 trials per condition. Each block lasted for about 15 minutes. The expected coefficient of variation for the threshold estimates is roughly 10% [48].

Participants sat upright on a chair placed on the MOOG platform in a completely dark room that eliminated the availability of any visual cues. The participant's body was secured to the chair using a five-point harness. Additionally, the participants' head was held firmly with the help of a helmet attached to the chair to prevent any movements of the head as the chair rotated. To minimize haptic cues caused by air motion, all the skin surfaces of the participants were fully covered (long sleeves, gloves and socks). External noise and other auditory cues were also reduced by using noise-cancelling earplugs and by playing white noise.

The direction of the motion was randomized across trials. In each motion trial participants were rotated either to the left or the right. They were instructed to indicate the perceived direction of motion by pushing either the left or the right button. For trials they were unsure of the direction, they were asked to make their best guess. Each motion stimuli was preceded by a "warning" tone and followed by a tone of higher pitch to mark the end of the trial. The subjects' responses to motion stimuli were recorded using a data acquisition system, which was used offline for data analysis.

As in previous studies, a three-down, one-up staircase adaptive algorithm was used to set the magnitude of the motion stimulus [9, 11, 49, 56, 85, 87]. "Three down" means that when the participant correctly reported the direction of three consecutive rotational trials, the velocity of the motion stimuli was reduced. "One up" means that when the participant incorrectly reported the direction of one trial the velocity of the stimuli was increased. We used the standard parameter estimation sequential testing algorithm (PEST) to determine the step size [86]. Testing for all participants started with large magnitude of rotations well above the typical yaw rotation threshold ($4^\circ/s$). The participants were given a mandatory break

between each block during which the lights were switched on and they were allowed to get up from the chair and stretch.

2.4 Threshold and Bias determination

Threshold and bias were determined separately for the two conditions (with and without vibration) by fitting a Gaussian cumulative distribution function to the data using a generalized linear model and probit link function [65, 67, 87]. Specifically, the function call `b=brglmfit(ω ,y,'binomial','link','probit')` in Matlab 2016b (The Mathworks, MA, USA) was used to obtain the fits, where ω is the vector containing amplitude of motion stimulus and y is the vector containing the participants' direction responses to the motion stimulus [12]. The output of this function contains elements that relate to the underlying Gaussian probability distribution, and were used to calculate threshold and bias. Threshold was defined as the standard deviation of the distribution and equal to the stimulus at which the subject correctly responds 84% of the times [41, 67]. Bias was defined as the mean of the distribution, located halfway between the lower and upper bounds of the psychometric function and equal to the stimulus that the subject correctly perceives as rightward 50% of the times. The sign of the bias denotes the direction along which the individual's perception is biased [67]. For example a bias equal to $-1.3^{\circ}/s$ denotes that on an average the individual perceives a rotation of magnitude equal to $1.3^{\circ}/s$ towards the left whenever exposed to null stimuli.

2.5 Statistical Analysis

All statistical tests were performed using SPSS V.20 (IBM Corp, Somers, NY). As in earlier studies, a log transformation was applied on the thresholds before performing any statistical tests [38, 49, 87]. Normality tests were performed before running any analyses. The distribution of both the biases and log-transformed thresholds of the two conditions were found to not vary significantly from normal.

Statistical testing was done using two approaches. First a paired t -test was used to compare thresholds and bias measured with and without vibration. Since a t -test can mask an effect when there is a substantial inter-subject variability in effect size, we also used a linear regression to determine the relationship between results obtained with and without vibration. A regression slope not significantly different from 1 was interpreted as a case in which there was no evidence of a difference in the measure between the two conditions across subjects. On the other hand, a slope significantly less than 1 was interpreted as a significant reduction in the measure with vibration from that without vibration. The slopes of the regression analyses were then tested using a one-sample t -test to determine if they were significantly different from a mean slope of 1.

3. Results

Fig. 2 shows perceptual responses and psychometric function fits to those responses for one subject. The dashed-dotted line indicates the level where the subject perceived 50% of the motions to be leftward and 50% of the motions to be rightward, which is referred to as the vestibular bias. The dashed line in the figure indicates the level where the subject perceived

84% of the motions to be rightward. The one-standard-deviation threshold is the difference in stimulus amplitude between the 50% and 84% levels (i.e., the 84% level after adjusting for vestibular bias). For this subject, bias was larger without vibration (Fig. 2A, dashed-dotted line) compared to with vibration (Fig. 2B, dashed-dotted line). The change in threshold for this subject was minimal.

Fig. 3A shows the relationship between thresholds measured with and without vibration, while Fig. 3B shows the relationship between biases measured with and without vibration. The paired *t*-test showed no significant differences in threshold or bias ($p>0.05$) measured with and without vibration.

A *t*-test can mask an effect when there is a substantial inter-subject variability in effect size. Thus, we also examined the effect of vibration on threshold using a linear regression between the log-transformed thresholds measured without vibration with those measured with vibration, which yielded a slope of 1.026 (S.E.= ± 0.065 , Fig. 3A, dashed line). This slope was not significantly different from a slope of 1 ($p=0.162$). This confirms that vibration had little effect on threshold. In contrast, when we examined the effect of vibration on bias using a linear regression between the biases with and without vibration, it yielded a slope of 0.4064 (S.E.= ± 0.109 , Fig. 3B, dashed line), which was found to be significantly less than a slope of 1 ($p<0.001$).

This result suggests that bias was significantly reduced in the presence of vibration when compared to that without vibration, but vibration had little effect on threshold. However, visual inspection of bias results indicated that one of the subjects might have been an outlier that could have disproportionately affected the regression slope; we now describe analyses that showed that this was not the case and that the subject had a minimal impact on the results. To objectively determine if there were outliers, we calculated the z-scores (standardized residuals) and Cook's distance. To find outliers using z-scores, the residuals of the data during model fitting were determined and standardized to a normal distribution, and any standardized residual with an absolute value >3.29 was flagged as an outlier [31]. Cook's distance measures the overall influence of a particular case over the parameters of the model, and any data with a Cook's distance >1 was flagged as an outlier [31]. None of the data points qualified as an outlier. Despite these results, we conducted a separate regression analysis without the visually-identified outlier to ensure that our result was not an artifact. Removal of the outlier changed the slope by a small amount (slope=0.331, S.E.= ± 0.210). In addition, we observed that some subjects had biases with different polarities with and without vibration. Since the interpretation of the effect of these subjects on the regression slope, as well as the physiological origins, are unclear, we also repeated analyses after excluding the three subjects whose biases switched polarity with vibration. This resulted in regression slope of 0.421 (S.E.= ± 0.119) with the outlier included and slope of 0.380 (S.E.= ± 0.240) without the outlier. These additional statistical tests provide further support that vibration changes bias by a factor of roughly 0.4.

4. Discussion

This study examined the effect of mastoid vibration on perceptual yaw rotation threshold and bias in healthy adults. The results show that vibration had little effect on yaw rotation threshold. There was some evidence that vibration reduced perceptual bias, although the results should be interpreted with caution. Analysis using linear regression found that there was a statistically significant reduction in bias, while analysis using paired *t*-test found no statistically significant change. Future testing with a larger sample size could potentially account for inter-subject variability and confirm the results obtained in this study. We note that *t*-tests may not detect an effect when there is substantial intersubject variability in effect size.

Evidence suggests that mastoid vibration broadly stimulates the vestibular organs, including the semicircular canals (SCC) and otolith organs, and that the relative activation of the two organs may depend on the specific stimuli. For example, E.D. Young et al. [93] recorded single unit activity of vestibular nerve fibers from adult squirrel monkeys in response to vibration applied to the frontal bone and found that SCC neuronal units are more sensitive to vibration than otolithic units and specifically most sensitive in the frequency range of 200–400 Hz. On the other hand, I.S. Curthoys et al. [19] reported that sinusoidal vibration (200–1,500 Hz) and short impulsive stimuli result in preferential activation of otolithic neurons as opposed to SCC neurons in guinea pigs. It has been proposed that the preferential tuning of otolithic afferents to rapid and sudden changes in accelerations facilitates the activation of these afferents to single clicks of vibratory stimulus [27]. Thus, our stimuli, consisting of frequencies in the range of 1–500 Hz, likely activated both the otolith organs and SCC. However, given the interspecies differences, there is a possibility that the otolith organs are more affected by vibration than the SCC. Future experiments could aim to compare these effects by testing translational perceptual thresholds (e.g., interaural) in addition to rotation. Emphasizing the importance of testing both, a recent study using electrical (galvanic) stochastic vestibular stimulation [33] found that while translation perceptual thresholds decreased in the presence of noise, SCC thresholds were unaffected.

We do not know the mechanism by which vibration could reduce vestibular bias. In fact, little is known about the physiologic basis for perceptual bias [35, 67]. It is speculated that perceptual vestibular bias is related to the imbalances in the peripheral or central vestibular function or a combination of both [74]. Given that vibration was applied anatomically close to the vestibular organs, it certainly is likely that peripheral factors play an important role in altering perceptual bias in this study although that does not rule out additional central contributions. One of the causes for peripheral imbalance might be the presence of structural asymmetries. For example, the orientation of SCC pairs is reported to be slightly different in right and left ears in certain mammals [4]. Similarly, asymmetries in the size and mass of otoconia have been found in invertebrates and these asymmetries at the peripheral level can have a direct effect on vestibular functions [3, 59, 60]. So, it could be possible that the structural asymmetries in the orientation of the SCC affects the fluid dynamics of the endolymph differently on the two sides, which in turn causes imbalance in the displacement of the hair cells and the activation of the vestibular neurons, resulting in perceptual bias. Bone conducted vibration is proposed to activate the vestibular receptors by modulating the

fluid pressure [64] and producing pressure waves that travel through the inner ear and displace the hair bundles [18]. It is possible that these fluid pressure waves somehow corrected for the differences in the fluid dynamics of the SCCs produced by the structural asymmetries and thus reduced perceptual bias in the current study.

We modeled six scenarios with different assumptions to determine how vestibular bias may arise from unequal neural activity on the left and right sides [67, 74], including additive (Fig. 4A–D) and multiplicative (Fig. 4E–H) asymmetries. We assumed that: a) perceived motion is determined by averaging bilateral activity, with the polarity of the left ear reversed, mimicking the standard push-pull SCC configuration in which the brain effectively inverts one signal during bilateral integration; b) perceptual bias arises from non-zero average activity with zero motion; c) neurons respond linearly with amplitude and direction, consistent with experimental data for small stimuli [26]; and d) perceptual asymmetries are not centrally compensated, since this is inconsistent with perceptual experimental results, despite evidence of central compensation in motor reflexes.

We assume a constant right-left difference in the additive case (Fig. 4A–D). As motion velocity becomes more positive, neural activity increases for the right SCC (dashed line) and decreases for the left SCC (solid line; polarity reversed for display purposes). Fig. 4B shows the predicted changes in responses if bilateral vibration increased activity by a constant amount only on the side with less baseline activity (i.e., the right side), resulting in average activity that is close to veridical and therefore consistent with our results. Despite evidence that vibration increases activity, we cannot explain how bilateral vibration would affect the less active side more. If both sides were equally affected (not shown), the two curves move in opposite directions by equal amounts, thus keeping the average the same. When correction occurs via an increase in sensitivity on the side with less baseline activity (Fig. 4C), i.e., a multiplicative correction, perceptual bias reduces at some amplitudes but increases overall. When correction occurs via equal bilateral reductions in sensitivity (Fig. 4D), perceptual bias is reduced. While this case is more plausible because vibration has bilateral effects, past experimental work found that vibration causes excitation rather than inhibition. We assume different right and left sensitivities for the multiplicative case (Fig. 4E). Since this predicts zero perceptual bias at zero motion amplitude, it does not match experimental results. Fig. 4F,G,H show the same correction models as for the additive case. In summary, Fig. 4B,D are consistent with our results, although not previous experimental evidence.

Although the vibratory stimulus used in the current study was meant to stimulate the vestibular organs peripherally, the factors that were used to generate the stimulus are generally used to provide tactile stimulation [21, 22, 54, 55, 75–77]. Although unlikely, it is possible that the vibration did not have a direct effect on the vestibular organs and the changes in bias that we observed are a consequence of tactile stimulation caused by vibration. Many behavioral studies have shown that vestibular-somatosensory interaction have important effects on perception, which are mainly due to anatomical overlap of neural sites that are involved in the processing of vestibular and somatosensory inputs. For example, improvement in perception of faint somatosensory stimuli was reported in the presence of caloric [28], galvanic [29] or natural vestibular stimulation (whole body

rotations) [30, 32]. The improvement in somatosensory perception in the presence of vestibular stimulation has been suggested to occur due to convergence of vestibular and somatosensory signals at bimodal neurons, where one sensory signal modulates the gain or synaptic connection in the afferent pathway of the other sensory signal. Several neurons that respond to both vestibular and somatosensory inputs have been identified by neuroimaging studies in certain regions of the brain [8, 95, 96]. Taken together, these findings suggest the possibility that changes in vestibular bias produced by vibration in this study could result from vibration (tactile stimulus) induced central modulation of the activity of the bimodal neuron that is responsible for compensating for any peripheral imbalances.

A few methodological limitations in this study are worth noting. Firstly, the broad vibration frequency range (1–500 Hz) used is known to stimulate a number of sensory organs that are proximate to the mastoid, many of which have known contributions to spatial orientation or multisensory interactions with vestibular cues. For example, the posterior and superior SCCs or otolith organs could have been activated, and although they are relatively insensitive to the yaw stimulus used, they interact in central spatial orientation mechanisms [2, 36, 66, 68]. Also, one of the cervical muscles (sternocleidomastoid muscle) could have been activated, which is known to affect postural responses [63], and thus may also play a role in self-motion perception. Furthermore, the auditory system was likely activated [25], and multisensory interactions occur between the auditory and vestibular systems in spatial processing [73, 88, 91]. In fact, auditory noise has been shown to change the processing of cross-modality cues [58]. Finally, cutaneous mechanoreceptors in the skin could have been activated [15, 43, 70, 77]. Additionally, the small sample size and limited vibratory and motion characteristics tested (yaw rotation, 1 Hz) are a few other limitations. Nonetheless, our statistical analyses, including the exclusion of artifacts from potential outliers, increase confidence in our conclusions. We expect that this investigation will provide the basis for the design of future experiments to address more specific questions using mechanical vibration like the most effective frequency range and intensity of vibration, and comparison of vibration effects on direction recognition thresholds of other motion planes (such as inter-aural translation or roll tilt).

Acknowledgment

This research was supported by NIH/NIDCD R03-DC013635. The authors thank Dr. Daniel M. Merfeld, the Director of the JVPL for providing us access to the MOOG device and Dr. Beom-Chan Lee for designing the software to precisely control factor activation.

References

- [1]. Aihara T, Kitajo K, Nozaki D and Yamamoto Y, How does stochastic resonance work within the human brain? – Psychophysics of internal and external noise, *Chemical Physics* 375 (2010), 616–624.
- [2]. Angelaki DE, McHenry MQ, Dickman JD, Newlands SD and Hess BJM, Computation of inertial motion: Neural strategies to resolve ambiguous otolith information, *Journal of Neuroscience* 19 (1999), 316–327. [PubMed: 9870961]
- [3]. Anken RH, Werner K, Ibsch M and Rahmann H, Fish inner ear otolith size and bilateral asymmetry during development, *Hear Res* 121 (1998), 77–83. [PubMed: 9682810]
- [4]. Berlin JC, Kirk EC and Rowe TB, Functional implications of ubiquitous semicircular canal non-orthogonality in mammals, *PLoS One* 8 (2013), e79585. [PubMed: 24260256]

- [5]. Bermudez Rey MC, Clark TK, Wang W, Leeder T, Bian Y and Merfeld DM, Vestibular Perceptual Thresholds Increase above the Age of 40, *Front Neurol* 7 (2016), 162. [PubMed: 27752252]
- [6]. Borah J, Young LR and Curry RE, Optimal estimator model for human spatial orientation, *Ann. N. Y. Acad. Sci* 545 (1988), 51–73. [PubMed: 3071213]
- [7]. Borel L and Ribot-Ciscar E, Improving postural control by applying mechanical noise to ankle muscle tendons, *Experimental Brain Research* 234 (2016), 2305–2314. [PubMed: 27021075]
- [8]. Bottini G, Sterzi R, Paulesu E, Vallar G, Cappa SF, Erminio F, Passingham RE, Frith CD and Frackowiak RS, Identification of the central vestibular projections in man: a positron emission tomography activation study, *Exp Brain Res* 99 (1994), 164–169. [PubMed: 7925790]
- [9]. Butler JS, Smith ST, Campos JL and Bulthoff HH, Bayesian integration of visual and vestibular signals for heading, *J Vis* 10 (2010), 23.
- [10]. Butler JS, Smith ST, Campos JL and Bulthoff HH, Bayesian integration of visual and vestibular signals for heading, *J. Vis* 10 (2010), 23.
- [11]. Chaudhuri SE, Karmali F and Merfeld DM, Whole body motion-detection tasks can yield much lower thresholds than direction-recognition tasks: implications for the role of vibration, *J Neurophysiol* 110 (2013), 2764–2772. [PubMed: 24068754]
- [12]. Chaudhuri SE and Merfeld DM, Signal detection theory and vestibular perception: III. Estimating unbiased fit parameters for psychometric functions, *Exp Brain Res* 225 (2013), 133–146. [PubMed: 23250442]
- [13]. Chien JH, Mukherjee M and Stergiou N, Mastoid Vibration Affects Dynamic Postural Control During Gait, *Ann Biomed Eng* 44 (2016), 2774–2784. [PubMed: 26833038]
- [14]. Collins JJ, Imhoff TT and Grigg P, Noise-enhanced tactile sensation, *Nature* 383 (1996), 770. [PubMed: 8893000]
- [15]. Collins JJ, Priplata AA, Gravelle DC, Niemi J, Harry J and Lipsitz LA, Noise-enhanced human sensorimotor function, *IEEE Eng Med Biol Mag* 22 (2003), 76–83. [PubMed: 12733463]
- [16]. Cousins S, Kaski D, Cutfield N, Seemungal B, Golding JF, Gresty M, Glasauer S and Bronstein AM, Vestibular Perception following Acute Unilateral Vestibular Lesions, *PLoS One* 8 (2013).
- [17]. Crane BT, Fore-aft translation aftereffects, *Exp. Brain Res* 219 (2012), 477–487. [PubMed: 22562589]
- [18]. Curthoys IS and Grant JW, How does high-frequency sound or vibration activate vestibular receptors?, *Exp Brain Res* 233 (2015), 691–699. [PubMed: 25567092]
- [19]. Curthoys IS, Kim J, McPhedran SK and Camp AJ, Bone conducted vibration selectively activates irregular primary otolithic vestibular neurons in the guinea pig, *Exp Brain Res* 175 (2006), 256–267. [PubMed: 16761136]
- [20]. Curthoys IS, Vulovic V, Burgess AM, Sokolic L and Goonetilleke SC, The response of guinea pig primary utricular and saccular irregular neurons to bone-conducted vibration (BCV) and air-conducted sound (ACS), *Hear Res* 331 (2016), 131–143. [PubMed: 26626360]
- [21]. Dettmer M, Pourmoghaddam A, Lee BC and Layne CS, Associations between Tactile Sensory Threshold and Postural Performance and Effects of Healthy Aging and Subthreshold Vibrotactile Stimulation on Postural Outcomes in a Simple Dual Task, *Curr Gerontol Geriatr Res* 2016 (2016), 9797369. [PubMed: 27143967]
- [22]. Dettmer M, Pourmoghaddam A, Lee BC and Layne CS, Do Aging and Tactile Noise Stimulation Affect Responses to Support Surface Translations in Healthy Adults?, *Curr Gerontol Geriatr Res* 2016 (2016), 2941964. [PubMed: 27195007]
- [23]. Diaz-Artiles A, Priesol AJ, Clark TK, Sherwood DP, Oman CM, Young LR and Karmali F, The Impact of Oral Promethazine on Human Whole-Body Motion Perceptual Thresholds, *J Assoc Res Otolaryngol* 18 (2017), 581–590. [PubMed: 28439720]
- [24]. Dumas G, Curthoys IS, Lion A, Perrin P and Schmerber S, The Skull Vibration-Induced Nystagmus Test of Vestibular Function-A Review, *Front Neurol* 8 (2017), 41. [PubMed: 28337171]
- [25]. Evans EF, The frequency response and other properties of single fibres in the guinea-pig cochlear nerve, *The Journal of Physiology* 226 (1972), 263–287. [PubMed: 5083170]

- [26]. Fernandez C and Goldberg JM, Physiology of peripheral neurons innervating semicircular canals of the squirrel monkey. II. Response to sinusoidal stimulation and dynamics of peripheral vestibular system, *J Neurophysiol* 34 (1971), 661–675. [PubMed: 5000363]
- [27]. Fernandez C and Goldberg JM, Physiology of peripheral neurons innervating otolith organs of the squirrel monkey. III. Response dynamics, *J Neurophysiol* 39 (1976), 996–1008. [PubMed: 824414]
- [28]. Ferre ER, Bottini G and Haggard P, Vestibular modulation of somatosensory perception, *Eur J Neurosci* 34 (2011), 1337–1344. [PubMed: 21978189]
- [29]. Ferre ER, Day BL, Bottini G and Haggard P, How the vestibular system interacts with somatosensory perception: a sham-controlled study with galvanic vestibular stimulation, *Neurosci Lett* 550 (2013), 35–40. [PubMed: 23827220]
- [30]. Ferre ER, Kaliuzhna M, Herbelin B, Haggard P and Blanke O, Vestibular-somatosensory interactions: effects of passive whole-body rotation on somatosensory detection, *PLoS One* 9 (2014), e86379. [PubMed: 24466064]
- [31]. Field A, *Discovering Statistics Using SPSS*, SAGE Publications, 2009.
- [32]. Figliozzi F, Guariglia P, Silvetti M, Siegler I and Doricchi F, Effects of vestibular rotatory accelerations on covert attentional orienting in vision and touch, *J Cogn Neurosci* 17 (2005), 1638–1651. [PubMed: 16269102]
- [33]. Galvan-Garza R, *Enhancement of Perception with the Application of Stochastic Vestibular Stimulation*, Massachusetts Institute of Technology, 2016.
- [34]. Galvan-Garza RC, Clark TK, Mulavara AP and Oman CM, Exhibition of stochastic resonance in vestibular tilt motion perception, *Brain Stimul* (2018).
- [35]. Garcia-Perez MA and Alcala-Quintana R, Shifts of the psychometric function: distinguishing bias from perceptual effects, *Q. J. Exp. Psychol. (Hove.)* 66 (2013), 319–337. [PubMed: 22950887]
- [36]. Glasauer S and Merfeld D, Modelling three-dimensional vestibular responses during complex motion stimulation, *Three-dimensional kinematics of eye, head and limb movements* (1997), 387–398.
- [37]. Goel R, Kofman I, Jeevarajan J, De Dios Y, Cohen HS, Bloomberg JJ and Mulavara AP, Using Low Levels of Stochastic Vestibular Stimulation to Improve Balance Function, *PLoS One* 10 (2015), e0136335. [PubMed: 26295807]
- [38]. Grabherr L, Nicoucar K, Mast FW and Merfeld DM, Vestibular thresholds for yaw rotation about an earth-vertical axis as a function of frequency, *Exp Brain Res* 186 (2008), 677–681. [PubMed: 18350283]
- [39]. Gravelle DC, Laughton CA, Dhruv NT, Katdare KD, Niemi JB, Lipsitz LA and Collins JJ, Noise-enhanced balance control in older adults, *Neuroreport* 13 (2002), 1853–1856. [PubMed: 12395078]
- [40]. Green DM and Swets JA, *Signal Detection Theory and Psychophysics*, Wiley, New York, 1966.
- [41]. Green DM and Swets JA, *Signal detection theory and psychophysics*, Wiley, New York, 1966.
- [42]. Hamann KF and Schuster EM, Vibration-induced nystagmus - A sign of unilateral vestibular deficit, *ORL J Otorhinolaryngol Relat Spec* 61 (1999), 74–79. [PubMed: 10095196]
- [43]. Hijmans JM, Geertzen JH, Zijlstra W, Hof AL and Postema K, Effects of vibrating insoles on standing balance in diabetic neuropathy, *J Rehabil Res Dev* 45 (2008), 1441–1449. [PubMed: 19319766]
- [44]. Huterer M and Cullen KE, Vestibuloocular reflex dynamics during high-frequency and high-acceleration rotations of the head on body in rhesus monkey, *J. Neurophysiol* 88 (2002), 13–28. [PubMed: 12091529]
- [45]. Karlberg M, Vibration-induced ocular torsion and nystagmus after unilateral vestibular deafferentation, *Brain* 126 (2003), 956–964. [PubMed: 12615651]
- [46]. Karlberg M, Aw ST, Black RA, Todd MJ, MacDougall HG and Halmagyi GM, Vibration-induced ocular torsion and nystagmus after unilateral vestibular deafferentation, *Brain* 126 (2003), 956–964. [PubMed: 12615651]

- [47]. Karmali F, Bermudez Rey MC, Clark TK, Wang W and Merfeld DM, Multivariate Analyses of Balance Test Performance, Vestibular Thresholds, and Age, *Front Neurol* 8 (2017), 578. [PubMed: 29167656]
- [48]. Karmali F, Chaudhuri SE, Yi Y and Merfeld DM, Determining thresholds using adaptive procedures and psychometric fits: evaluating efficiency using theory, simulations, and human experiments, *Exp Brain Res* 234 (2016), 773–789. [PubMed: 26645306]
- [49]. Karmali F, Lim K and Merfeld DM, Visual and vestibular perceptual thresholds each demonstrate better precision at specific frequencies and also exhibit optimal integration, *J Neurophysiol* 111 (2014), 2393–2403. [PubMed: 24371292]
- [50]. Karmali F and Merfeld DM, A distributed, dynamic, parallel computational model: the role of noise in velocity storage, *J Neurophysiol* 108 (2012), 390–405. [PubMed: 22514288]
- [51]. Karmali F, Whitman GT and Lewis RF, Bayesian optimal adaptation explains age-related human sensorimotor changes, *J Neurophysiol* 119 (2018), 509–520. [PubMed: 29118202]
- [52]. Keywan A, Wuehr M, Pradhan C and Jahn K, Noisy Galvanic Stimulation Improves Roll-Tilt Vestibular Perception in Healthy Subjects, *Front Neurol* 9 (2018), 83. [PubMed: 29545766]
- [53]. Laurens J and Droulez J, Bayesian Processing of Vestibular Information, *Biological Cybernetics* 96 (2007), 389–404. [PubMed: 17146661]
- [54]. Lee BC, Martin BJ and Sienko KH, Directional postural responses induced by vibrotactile stimulations applied to the torso, *Exp Brain Res* 222 (2012), 471–482. [PubMed: 22968737]
- [55]. Lee BC, Martin BJ and Sienko KH, The effects of actuator selection on non-volitional postural responses to torso-based vibrotactile stimulation, *J Neuroeng Rehabil* 10 (2013), 21. [PubMed: 23406013]
- [56]. Leek MR, Adaptive procedures in psychophysical research, *Percept Psychophys* 63 (2001), 1279–1292. [PubMed: 11800457]
- [57]. Lewis RF, Priesol AJ, Nicoucar K, Lim K and Merfeld DM, Abnormal motion perception in vestibular migraine, *Laryngoscope* 121 (2011), 1124–1125. [PubMed: 21520135]
- [58]. Lugo E, Doti R and Faubert J, Ubiquitous crossmodal Stochastic Resonance in humans: auditory noise facilitates tactile, visual and proprioceptive sensations, *PLoS One* 3 (2008), e2860. [PubMed: 18682745]
- [59]. Lychakov DV, [Lateralization of behavioral reactions and otolith asymmetry], *Zh Evol Biokhim Fiziol* 49 (2013), 299–312. [PubMed: 24459865]
- [60]. Lychakov DV and Rebane YT, Fish otolith mass asymmetry: morphometry and influence on acoustic functionality, *Hear Res* 201 (2005), 55–69. [PubMed: 15721561]
- [61]. MacNeilage PR, Ganesan N and Angelaki DE, Computational Approaches to Spatial Orientation: From Transfer Functions to Dynamic Bayesian Inference, *Journal of Neurophysiology* 100 (2008), 2981–2996. [PubMed: 18842952]
- [62]. Magalhães FH and Kohn AF, Effectiveness of electrical noise in reducing postural sway: a comparison between imperceptible stimulation applied to the anterior and to the posterior leg muscles, *European Journal of Applied Physiology* 114 (2014), 1129–1141. [PubMed: 24563056]
- [63]. Magnusson M, Andersson G, Gomez S, Johansson R, Martensson A, Karlberg M and Fransson PA, Cervical muscle afferents play a dominant role over vestibular afferents during bilateral vibration of neck muscles, *J Vestib Res* 16 (2006), 127–136. [PubMed: 17312340]
- [64]. Maxwell AK, Banakis Hartl RM, Greene NT, Benichoux V, Mattingly JK, Cass SP and Tollin DJ, Semicircular Canal Pressure Changes During High-intensity Acoustic Stimulation, *Otol Neurotol* 38 (2017), 1043–1051. [PubMed: 28570420]
- [65]. McCullagh P and Nelder JA, *Generalized linear models*, Chapman and Hall, London; New York, 1989.
- [66]. Merfeld DM, Modeling the vestibule-ocular reflex of the squirrel monkey during eccentric rotation and roll tilt, *Experimental Brain Research* 106 (1995), 123–134. [PubMed: 8542968]
- [67]. Merfeld DM, Signal detection theory and vestibular thresholds: I. Basic theory and practical considerations, *Exp Brain Res* 210 (2011), 389–405. [PubMed: 21359662]
- [68]. Merfeld DM and Zupan LH, Neural processing of gravito-inertial cues in humans. III. Modeling tilt and translation responses, *Journal of Neurophysiology* 87 (2002), 819–833. [PubMed: 11826049]

- [69]. Moss F, Ward LM and Sannita WG, Stochastic resonance and sensory information processing: a tutorial and review of application, *Clin Neurophysiol* 115 (2004), 267–281. [PubMed: 14744566]
- [70]. Mountcastle VB, LaMotte RH and Carli G, Detection thresholds for stimuli in humans and monkeys: comparison with threshold events in mechanoreceptive afferent nerve fibers innervating the monkey hand, *J Neurophysiol* 35 (1972), 122–136. [PubMed: 4621505]
- [71]. Mulavara AP, Fiedler MJ, Kofman IS, Wood SJ, Serrador JM, Peters B, Cohen HS, Reschke MF and Bloomberg JJ, Improving balance function using vestibular stochastic resonance: optimizing stimulus characteristics, *Exp Brain Res* 210 (2011), 303–312. [PubMed: 21442221]
- [72]. Pal S, Rosengren SM and Colebatch JG, Stochastic galvanic vestibular stimulation produces a small reduction in sway in Parkinson's disease, *J Vestib Res* 19 (2009), 137–142. [PubMed: 20448339]
- [73]. Perrott DR, Elfner LF and Homick JL, Auditory spatial orientation, *Perception & Psychophysics* 5 (1969), 189–192.
- [74]. Peterka RJ and Benolken MS, Relation between perception of vertical axis rotation and vestibulo-ocular reflex symmetry, *J Vestib Res* 2 (1992), 59–69. [PubMed: 1342384]
- [75]. Priplata A, Niemi J, Salen M, Harry J, Lipsitz LA and Collins JJ, Noise-enhanced human balance control, *Phys Rev Lett* 89 (2002), 238101. [PubMed: 12485044]
- [76]. Priplata AA, Niemi JB, Harry JD, Lipsitz LA and Collins JJ, Vibrating insoles and balance control in elderly people, *Lancet* 362 (2003), 1123–1124. [PubMed: 14550702]
- [77]. Priplata AA, Patriiti BL, Niemi JB, Hughes R, Gravelle DC, Lipsitz LA, Veves A, Stein J, Bonato P and Collins JJ, Noise-enhanced balance control in patients with diabetes and patients with stroke, *Ann Neurol* 59 (2006), 4–12. [PubMed: 16287079]
- [78]. Ramachandran R and Lisberger SG, Normal performance and expression of learning in the vestibulo-ocular reflex (VOR) at high frequencies, *J. Neurophysiol* 93 (2005), 2028–2038. [PubMed: 15548626]
- [79]. Ribot-Ciscar E, Hospod V and Aimonetti JM, Noise-enhanced kinaesthesia: a psychophysical and microneurographic study, *Exp Brain Res* 228 (2013), 503–511. [PubMed: 23712687]
- [80]. Roditi RE and Crane BT, Directional asymmetries and age effects in human self-motion perception, *J. Assoc. Res. Otolaryngol* 13 (2012), 381–401. [PubMed: 22402987]
- [81]. Saginaw MA, *Eye Movement Studies with a Vestibular Prosthesis*, Massachusetts Institute of Technology, 2010.
- [82]. Saginaw MA, Gong W, Haburcakova C and Merfeld DM, Eye movements above 100 Hz studied with vestibular electrical prosthesis, 2007.
- [83]. Sasaki H, Todorokihara M, Ishida T, Miyachi J, Kitamura T and Aoki R, Effect of noise on the contrast detection threshold in visual perception, *Neurosci Lett* 408 (2006), 94–97. [PubMed: 16996210]
- [84]. Scinicariello AP, Inglis JT and Collins JJ, The effects of stochastic monopolar galvanic vestibular stimulation on human postural sway, *J Vestib Res* 12 (2002), 77–85. [PubMed: 12867666]
- [85]. Soyka F, Robuffo Giordano P, Beykirch K and Bulthoff HH, Predicting direction detection thresholds for arbitrary translational acceleration profiles in the horizontal plane, *Exp Brain Res* 209 (2011), 95–107. [PubMed: 21234751]
- [86]. Taylor MM and Creelman CD, PEST: Efficient Estimates on Probability Functions, *The Journal of the Acoustical Society of America* 41 (1967), 782–787.
- [87]. Valko Y, Lewis RF, Priesol AJ and Merfeld DM, Vestibular labyrinth contributions to human whole-body motion discrimination, *J Neurosci* 32 (2012), 13537–13542. [PubMed: 23015443]
- [88]. Voss P, Auditory Spatial Perception without Vision, *Frontiers in Psychology* 7 (2016).
- [89]. Watamaniuk SN and Heinen SJ, Human smooth pursuit direction discrimination, *Vision Res* 39 (1999), 59–70. [PubMed: 10211396]
- [90]. Welgampola MS, Rosengren SM, Halmagyi GM and Colebatch JG, Vestibular activation by bone conducted sound, *J Neurol Neurosurg Psychiatry* 74 (2003), 771–778. [PubMed: 12754349]
- [91]. Wigderson E, Nelken I and Yarom Y, Early multisensory integration of self and source motion in the auditory system, *Proceedings of the National Academy of Sciences* 113 (2016), 8308–8313.

- [92]. Xie S, Guo J, Wu Z, Qiang D, Huang J, Zheng Y, Yao Q, Chen S and Tian D, Vibration-induced nystagmus in patients with unilateral peripheral vestibular disorders, *Indian J Otolaryngol Head Neck Surg* 65 (2013), 333–338.
- [93]. Young ED, Fernandez C and Goldberg JM, Responses of squirrel monkey vestibular neurons to audio-frequency sound and head vibration, *Acta Otolaryngol* 84 (1977), 352–360. [PubMed: 303426]
- [94]. Zeng FG, Fu QJ and Morse R, Human hearing enhanced by noise, *Brain Res* 869 (2000), 251–255. [PubMed: 10865084]
- [95]. zu Eulenburg P, Baumgartner U, Treede RD and Dieterich M, Interoceptive and multimodal functions of the operculo-insular cortex: tactile, nociceptive and vestibular representations, *Neuroimage* 83 (2013), 75–86. [PubMed: 23800791]
- [96]. zu Eulenburg P, Caspers S, Roski C and Eickhoff SB, Meta-analytical definition and functional connectivity of the human vestibular cortex, *Neuroimage* 60 (2012), 162–169. [PubMed: 22209784]
- [97]. Zupan LH and Merfeld DM, Interaural self-motion linear velocity thresholds are shifted by roll vection, *Exp. Brain Res* 191 (2008), 505–511. [PubMed: 18843487]




Fig. 1: An example of one cycle of sinusoidal acceleration of yaw rotation. This corresponds to cosine bell velocity and sigmoidal displacement profiles at 1 Hz motion with a displacement of 1 degree.




Fig. 2:
Example of typical subject's responses fit with Gaussian cumulative distribution psychometric functions (A) without vibration; B) with vibration. The position of the circles represents the fraction rightward out of all responses at given amplitude and the size of the circle indicates the total number of responses. In some cases, responses for stimuli within smaller intervals were pooled for display only.




Fig. 3:

A) Relationship between thresholds with and without vibration. B) Relationship between bias with and without vibration. The dashed black line represents the regression line and the circular dots represent each subject's experimentally-measured thresholds and biases. The solid line (unity line) represents a line with slope equal to 1. The area enclosed within the shaded grey region represents the 95% confidence interval of the regression line. The data point on the far left of figure B represents the visually identified outlier. The red colored dot represents the data of a subject who had difficulty in performing the task. Both the data points were used in all analyses since they were objectively not found to be outliers.




Fig. 4:

A) The neural representations of the left (solid line) and right (dashed line) semicircular canals as a function of motion stimulus amplitude for the additive asymmetry case. The neural activity of the right side is assumed to be less than the left side by a constant amount. B,C,D) As described in the text, three scenarios are considered for how the activity shown in (A) could be modified by vibration. E) In the multiplicative case, the sensitivity of the right side is assumed to be lesser than the left side. F,G,H) As described in the text, three scenarios are considered for how the activity shown in (D) could be modified by vibration.